


# Tuning the Optical Properties of Silicon Quantum Dots via Surface Functionalization with Conjugated Aromatic Fluorophores

Mohammed Abdelhameed,<sup>a</sup> Diego Rota Martira,<sup>b</sup> Shalimar Chen,<sup>c</sup> William Z. Xu,<sup>a</sup> Olabode O.


Oyeneye,<sup>a</sup> Subrata Chakrabarti,<sup>c</sup> Eli Zysman-Colman<sup>b</sup> and Paul A. Charpentier<sup>a</sup>

<b>Figure S1.</b> FTIR spectra of 3-Ethynylperylene, 9-Ethynylphenanthrene, and 1-Ethynylpyrene. ....	<b>3</b>
<b>Figure S2.</b> XPS spectra of C 1S for SQD-pyrene. ....	<b>4</b>
<b>Figure S3.</b> UV-Vis absorption, excitation, and emission spectra of SQD-heptene collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 360$ nm). ....	<b>5</b>
<b>Figure S4.</b> UV-Vis absorption, excitation, and emission spectra of SQD-heptene collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 315$ nm). ....	<b>6</b>
<b>Figure S5.</b> UV-Vis spectra, excitation, and emission spectra of Perylene and SQD-perylene collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 360$ nm). ....	<b>7</b>
<b>Figure S6.</b> UV-Vis spectra, Excitation, and Emission spectra of Pyrene and SQD-pyrene collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 360$ nm). ....	<b>8</b>
<b>Figure S7.</b> UV-Vis spectra, excitation, and emission spectra of Phenanthrene and SQD-phenanthrene collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 315$ nm). ....	<b>9</b>
<b>Figure S8.</b> Emission spectra of SQD-perylene collected in degassed DCM at 298 K upon photoexcitation at 360 nm and at 450 nm. ....	<b>10</b>
<b>Figure S9.</b> Emission spectra of SQD-pyrene collected in degassed DCM at 298 K upon photoexcitation at 360 nm and at 348 nm. ....	<b>11</b>
<b>Figure S10.</b> Emission spectra of SQD-phenanthrene collected in degassed DCM at 298 K upon photoexcitation at 360 nm and at 315 nm. ....	<b>12</b>
<b>Figure S11.</b> Emission spectra of SQD-pyrene collected in degassed DCM at 298 K at a concentration of 100 $\mu\text{M}$ , 50 $\mu\text{M}$ , 10 $\mu\text{M}$ and 1 $\mu\text{M}$ ( $\lambda_{\text{ex}} = 360$ nm). ....	<b>13</b>
<b>Figure S12.</b> Emission spectra of Pyrene collected in degassed DCM at 298 K at a concentration of 1.5 mM, 550 $\mu\text{M}$ , 368 $\mu\text{M}$ and 176 $\mu\text{M}$ ( $\lambda_{\text{ex}} = 360$ nm). ....	<b>14</b>
<b>Figure S13.</b> Emission decay of SQD-heptene monitored at 448 nm collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 378$ nm). ....	<b>14</b>
<b>Figure S14.</b> Emission decay of perylene monitored at 467 nm collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 378$ nm). ....	<b>15</b>
<b>Figure S15.</b> Emission decay of SQD-perylene monitored at 519 nm collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 378$ nm). ....	<b>15</b>
<b>Figure S16.</b> Emission decay of pyrene monitored at 384 nm collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 378$ nm). ....	<b>16</b>
<b>Figure S17.</b> Emission decay of SQD-pyrene monitored at 394 nm collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 378$ nm). ....	<b>16</b>


<b>Figure S18.</b> Emission decay of phenanthrene monitored at 376 nm collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 378$ nm). .....	<b>17</b>
<b>Figure S19.</b> Emission decay of SQD-phenanthrene monitored at 446 nm collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 378$ nm). .....	<b>17</b>
<b>Figure S20.</b> Emission spectra of Perylene, Pyrene, Phenanthrene, SQD-perylene, SQD-pyrene, and SQD-phenanthrene. ....	<b>18</b>
<b>Figure S21.</b> Emission spectra of SQD-perylene (blue) and SQD-pyrene (orange) in water.....	<b>19</b>
<b>Figure S22.</b> Confocal images of HeLa cells for DIC images (panels A and D), fluorescence images (panels B and E), and merged images (panels C and F).....	<b>20</b>




**Figure S1.** FTIR spectra of 3-Ethynylperylene (a), 9-Ethynylphenanthrene (b), and 1-Ethynylpyrene (c).




**Figure S2.** XPS spectra of C 1S for SQD-pyrene




**Figure S3.** UV-Vis absorption, excitation ( $\lambda_{\text{em}} = 448$  nm), and emission spectra of SQD-heptene collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 360$  nm).



**Figure S4.** UV-Vis absorption, excitation ( $\lambda_{\text{em}} = 374$  nm), and emission spectra of SQD-heptene collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 315$  nm).




**Figure S5.** UV-Vis spectra of perylene (dotted orange line) and SQD-perylene (solid orange line); excitation spectra of perylene (dotted green line,  $\lambda_{\text{em}} = 467$  nm) and SQD-perylene (solid green line,  $\lambda_{\text{em}} = 518$  nm); and emission spectra of perylene (dotted red line) and SQD-perylene (solid red line) collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 360$  nm).




**Figure S6.** UV-Vis spectra of pyrene (dotted orange line) and SQD-pyrene (solid orange line); excitation spectra of pyrene (dotted green line,  $\lambda_{\text{em}} = 384$  nm) and SQD-pyrene (solid green line,  $\lambda_{\text{em}} = 397$  nm); and emission spectra of pyrene (dotted red line) and SQD-pyrene (solid red line) collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 360$  nm).







**Figure S7.** UV-Vis spectra of phenanthrene (dotted orange line) and SQD-phenanthrene (solid orange line); excitation spectra of Phenanthrene (dotted green line,  $\lambda_{\text{em}} = 377$  nm) and SQD-phenanthrene (solid green line,  $\lambda_{\text{em}} = 447$  nm); and emission spectra of phenanthrene (dotted red line) and SQD-phenanthrene (solid red line) collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 315$  nm).




**Figure S8.** Emission spectra of SQD-perylene collected in degassed DCM at 298 K upon photoexcitation at 360 nm (light-blue line) and at 450 nm (red line).




**Figure S9.** Emission spectra of SQD-pyrene collected in degassed DCM at 298 K upon photoexcitation at 360 nm (light-blue line) and at 348 nm (red line).




**Figure S10.** Emission spectra of SQD-phenanthrene collected in degassed DCM at 298 K upon photoexcitation at 360 nm (light-blue line) and at 315 nm (red line).




**Figure S11.** Emission spectra of SQD-pyrene collected in degassed DCM at 298 K at a concentration of 100  $\mu$ M (red line), 50  $\mu$ M (purple line), 10  $\mu$ M (orange line) and 1  $\mu$ M (blue line) ( $\lambda_{\text{ex}} = 360$  nm).




**Figure S12.** Emission spectra of pyrene collected in degassed DCM at 298 K at a concentration of 1.5 mM (blue line), 550  $\mu\text{M}$  (orange line), 368  $\mu\text{M}$  (green line) and 176  $\mu\text{M}$  (red line) ( $\lambda_{\text{ex}} = 360$  nm).




**Figure S13.** Emission decay of SQD-heptene monitored at 448 nm collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 378$  nm).




**Figure S14.** Emission decay of perylene monitored at 467 nm collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 378$  nm).



**Figure S15.** Emission decay of SQD-perylene monitored at 519 nm collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 378$  nm).




**Figure S16.** Emission decay of pyrene monitored at 384 nm collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 378$  nm).




**Figure S17.** Emission decay of SQD-pyrene monitored at 394 nm collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 378$  nm).







**Figure S18.** Emission decay of phenanthrene monitored at 376 nm collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 378$  nm).



**Figure S19.** Emission decay of SQD-phenanthrene monitored at 446 nm collected in degassed DCM at 298 K ( $\lambda_{\text{ex}} = 378$  nm).




**Figure S20.** Emission spectra of perylene ( $\lambda_{\text{ex}} = 360$  nm, dotted red line) and SQD-perylene ( $\lambda_{\text{ex}} = 360$  nm, solid red line); pyrene ( $\lambda_{\text{ex}} = 360$  nm, dotted blue line) and SQD-pyrene ( $\lambda_{\text{ex}} = 360$  nm, solid blue-line); phenanthrene ( $\lambda_{\text{ex}} = 315$  nm, dotted yellow line) and SQD-phenanthrene ( $\lambda_{\text{ex}} = 315$  nm, solid yellow line) collected in degassed DCM at 298 K.



**Figure S21.** Emission spectra of SQD-perylene ( $\lambda_{\text{ex}} = 440$  nm, blue) and SQD-pyrene ( $\lambda_{\text{ex}} = 350$  nm, orange) in water.

### Fluorescent cellular imaging study

In figure 5, we utilized a confocal microscope to collect the photoluminescence obtained from HeLa cells that were incubated with SQD-perylene and SQD-pyrene. Cells imaging was collected at two different channels of 400-450 and 500-550 nm in the presence of a combination of short- and long-pass filters, and at an excitation wavelength of 405 nm. When the emission was monitored using the 440-450 nm channel, a blue emission was obtained as shown in figure 5 while other emission wavelengths were filtered out. On the other hand, figure S22 below shows the green emission collected when the 500-550 nm channel was utilized. The broad emission spectrum of both SQD-perylene and SQD-pyrene has enabled to collect blue and green emission color when the channels 400-450 and 500-550 nm were utilized, respectively.



**Figure S22.** Confocal images of HeLa cells for DIC images (panels A and D), fluorescence images (panels B and E), and merged images (panels C and F).